


CZY  
TACHIONY  
ZBAWIĄ TEORIĘ  
KWANTOWĄ?


ANDRZEJ DRAGAN  
UNIWERSYTET WARSZAWSKI  
NATIONAL UNIVERSITY OF SINGAPORE




ANDRZEJ DĄBROWSKI  
UNIWERSYTET WARSZAWSKI  
NATIONAL UNIVERSITY OF SINGAPORE

1057829,8 (3,2) kHz


1057833,8 (6B) kHz






One might still like to ask: "How does it work? What is the machinery behind the law?" No one has found any machinery behind the law. No one can "explain" any more than we have just "explained." No one will give you any deeper representation of the situation. We have no ideas about a more basic mechanism from which these results can be deduced.


Challenge accepted!



One might still like to ask: "How does it work?" No one has found any machinery behind the law. No one can "explain" any more. We have just "explained." No one will give you any deeper representation of the situation. We have no idea what the basic mechanism from which these results








BORN  
BONDI  
FEYNMAN  
PAULI  
ROHLICH  
PEIERLS  
BOULWARE

~~GRV~~

BORN  
BONDI  
FEYNMAN  
PAULI  
ROHLICH  
PEIERLS  
BOULWARE



What I cannot create,  
I do not understand.

W/M/const x soft. P.

I know how to solve every  
problem that has been solved

TO LEARN:

Bethe Ansatz Probs.

Kondo  $\rightarrow$

2-D Hall,

accel. Temp

Non linear Derivat. Hydro

$$\textcircled{A} \quad f = u(k, a)$$

$$g = 4(t, z) u(k, z)$$

$$\textcircled{B} \quad f = 2|k, a| (u, a)$$

What I cannot create,  
I do not understand.

W/M/const x soft. P.

I know how to solve every  
problem that has been solved

TO LEARN:

Bethe Ansatz Probs.

Kondo ↑

J.B. Halle

accel. Temp

Non linear Derivat. Hybro


$$\textcircled{A} \quad f = u(k, a)$$

$$g = 4(t, z) u(k, z)$$

$$\textcircled{B} \quad f = 2|k, a| (u, a)$$



$$| \uparrow \uparrow \rangle + | \downarrow \downarrow \rangle$$



$$\left\{ \begin{array}{l} x' = \frac{x - vt}{\sqrt{1 - v^2/c^2}} \\ t' = \frac{t - vx/c^2}{\sqrt{1 - v^2/c^2}} \end{array} \right.$$

日本

LED  
ON

$$\left\{ \begin{array}{l} x' = \frac{x - vt}{\sqrt{1 - v^2/c^2}} \\ t' = \frac{t - vx/c^2}{\sqrt{1 - v^2/c^2}} \end{array} \right.$$




$$x' = \frac{x - vt}{\sqrt{1 - v^2/c^2}}$$


$$t' = \frac{t - v/c^2 x}{\sqrt{1 - v^2/c^2}}$$

(CZAS)

(PREZESTRZEN)




$$\left\{ \begin{array}{l} x' = \frac{x - vt}{\sqrt{1 - v^2/c^2}} \\ t' = \frac{t - v/c^2 x}{\sqrt{1 - v^2/c^2}} \end{array} \right.$$



$$2 \cdot (2AS) + \sqrt{3} \cdot (\text{PRZESTRZEN}) \\ = (CZAS)$$

$$\left\{ \begin{array}{l} x' = \frac{x - vt}{\sqrt{1 - v^2/c^2}} \\ t' = \frac{t - v/c^2 x}{\sqrt{1 - v^2/c^2}} \end{array} \right.$$



2 · (CZAS) +  $\sqrt{3} \cdot (\text{PRZESTRZEN})$


$$= (\cancel{\text{CZAS}})$$

$$= (\text{PRZESTRZEN})$$


$$\left\{ \begin{array}{l} x' = -\frac{v}{n} \frac{x-vt}{\sqrt{\frac{v^2}{c^2}-1}} \\ t' = -\frac{v}{n} \frac{t-\frac{v}{c^2}x}{\sqrt{\frac{v^2}{c^2}-1}} \end{array} \right.$$

$$\left\{ \begin{array}{l} x' = -\frac{v}{N} \frac{x-vt}{\sqrt{v^2/c^2-1}} \\ t' = -\frac{v}{N} \frac{t-v/c^2 x}{\sqrt{v^2/c^2-1}} \end{array} \right.$$

$$\boxed{v > c}$$



$$v \rightarrow \infty \Rightarrow \left\{ \begin{array}{l} x' = ct \\ ct' = x \end{array} \right.$$



# New Journal of Physics

The open access journal at the forefront of physics

PAPER • OPEN ACCESS

# Quantum principle of relativity

Andrzej Dragan<sup>1,2</sup> and Artur Ekert<sup>2,3</sup>

<sup>1</sup> Institute of Theoretical Physics, University of Warsaw

<sup>2</sup> Centre for Quantum Technologies, National University of Singapore


<sup>3</sup> Mathematical Institute, University of Oxford

Deutsche Physikalische Gesellschaft





IOP Institute of Physics


69247 Total downloads

























LOKALNA, DETERMINISTYCZNA  
I RELATYWISTYCZNA TEORIA  
NIEMOŻLIWA


$c t$  $x$  $c t'$  $x'$ 

$$c t' = x$$


$$x' = c t$$








SUPERPOSITION  
NIEUNIKIONE



$$P(A \rightarrow B) = \frac{1}{n_B} |e^{i\phi_1} + \dots + e^{i\phi_n}|^2$$

CZY TACHIONY  
ISTNIEJĄ ?



PRÓZNA

CZY TACHIONY  
ISTNIEJĄ ?

